

CHITOSAN MICROPARTICLES ORAL DELIVERY SYSTEM OF RELAXING PEPTIDES

Patrícia Batista,^[a] Pedro Castro,^[a] Ana Raquel Madureira,^[a] Bruno Sarmiento,^[b,c] Manuela Pintado^[a]

[a] Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina – Laboratório Associado, Escola Superior de Biotecnologia, Rua Arquiteto Lobão Vital, 172, 4200-374 Porto, Portugal

[b] CESPU, Instituto de Investigação e Formação Avançada em Ciências e Tecnologias da Saúde, Rua Central de Gandra 1317, 4585-116 Gandra, Portugal

[c] i3S - Instituto de Investigação e Inovação em Saúde/INEB - Instituto Nacional de Engenharia Biomédica, Universidade do Porto, Rua Alfredo Allen 208, 4200-393 Porto, Portugal

Bioactive peptides can prevent or control diseases and exert a variety of activities including relaxing function. Microencapsulation of bioactive peptides may constitute a valuable process for preserving their integrity during processing and digestion [1, 2]. Chitosan is a prominent natural biomaterial that is used in delivery systems [3-6]. In the present study, chitosan microparticles loaded with a relaxing peptide were prepared by ionic gelation, with a mean size of $1.813 \pm 0.802 \mu\text{m}$, zeta potential $43.49 \pm 9.09 \text{ mV}$ and polydispersion index 0.89. This delivery system was characterized in terms of particle size analysis, Fourier Transform Infrared, peptide encapsulation efficiency, bioavailability and cytotoxicity through *in vitro* assays. The results demonstrated that ca. 86% of the peptide was loaded into the chitosan microparticles. The microparticles showed to be biocompatible. Cell viability was measured by MTT assay and showed an absence of cell toxicity following exposure into formulation contact. Peptide-loaded chitosan microparticles and peptide-loaded chitosan microparticles into guar-gum films showed a faster and distinct permeability than a free peptide. So, peptide-loaded chitosan microparticles and peptide-loaded chitosan microparticles into guar-gum films might be a promising carrier for relaxing peptide buccal administration.

References

- [1] P. M. Castro, F. Sousa, R. Magalhães, V. M. P. Ruiz-Henestrosa, A. M. Pilosof, A. R. Madureira, B. Sarmiento and M. E. Pintado, Incorporation of beads into oral films for buccal and oral delivery of bioactive molecules, *Carbohydr Polym.* **2018** *194* 411-421.
- [2] D. J. McClements, Encapsulation, protection, and delivery of bioactive proteins and peptides using nanoparticle and microparticle systems: A review, *Adv. Colloid Interface Sci.* **2018** *253* 1-22.
- [3] S. M. Ahsan, M. Thomas, K. K. Reddy, S. G. Sooraparaju, A. Asthana and I. Bhatnagar, Chitosan as biomaterial in drug delivery and tissue engineering, *Int J Biol Macromol.* **2017** *110* 97-109.
- [4] D.-H. Ngo, T.-S. Vo, D.-N. Ngo, K.-H. Kang, J.-Y. Je, H. N.-D. Pham, H.-G. Byun and S.-K. Kim, Biological effects of chitosan and its derivatives, *Food Hydrocolloids* **2015** *51* 200-216.
- [5] R. Khlibsuwan, F. Siepmann, J. Siepmann and T. Pongjanyakul, Chitosan-clay nanocomposite microparticles for controlled drug delivery: Effects of the MAS content and TPP crosslinking, *J Drug Deliv Sci Technol* **2017** *40* 1-10.
- [6] L. B. Rodriguez, A. Avalos, N. Chiaia and A. Nadarajah, Effect of formulation and process parameters on chitosan microparticles prepared by an emulsion crosslinking technique, *AAPS PharmSciTech* **2017** *18*(4) 1084-1094.