



SOCIEDADE
PORTUGUESA
DE QUÍMICA

ANALÍTICA 2018


9th Meeting of Division of Analytical Chemistry

26-27 MARCH, PORTO-PORTUGAL

FFUP/ICBAS – UNIVERSITY OF PORTO

www.analitica2018.eventos.chemistry.pt

**BOOK OF
ABSTRACTS**



POLYMER INCLUSION MEMBRANES (PIMS) AS AN ALTERNATIVE FOR ON-LINE SOLID PHASE EXTRACTION (SPE) IN FLOW ANALYSIS

Tânia C. F. Ribas¹, Charles F. Croft², Raquel B. R. Mesquita¹, M. Inês G. S. Almeida², Spas D. Kolev², António O. S. S. Rangel¹

¹*Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina - Laboratório Associado, Escola Superior de Biotecnologia, Porto, Portugal*

²*School of Chemistry, The University of Melbourne, VIC, 3010, Australia
tpedro@porto.ucp.pt*

A membrane can be considered as a selective barrier between two phases; in recent years, membrane-based processes have been subject of high interest in many fields, namely biotechnology, wastewater treatment and others [1]. Membrane research has attracted considerable efforts in recent years to provide a better understanding of membrane-based separation and improve its efficiency. This line of investigation has led to the development of polymer inclusion membranes (PIMs) which are thin, flexible and stable polymeric films that can selectively separate chemical species of interest [2]. PIMs are fabricated by solvent casting a solution containing an extractant (the selective agent) and a base polymer that stably encapsulates the extractant. Some PIMs may also contain a plasticizer and/or chemical modifiers.

Solid phase extraction (SPE) is a widely used technique applied to the sample pre-treatment in analytical chemistry. This technique is associated to a significant number of advantages as it can selectively separate the analyte, or it could remove matrix interferences as a clean-up step in complex matrices. In certain conditions, SPE could also be a strategy for pre-concentration of a specific analyte present in low concentrations.

In this work, PIMs fabricated from 45 wt% di-(2-ethylhexyl)phosphoric acid (D2EHPA) and 55 wt% PVC are explored as an alternative sorbent material for SPE aiming for the separation and/or pre-concentration of Zn(II). This work aims to show that this novel SPE method can be used to separate and/or pre-concentrate Zn(II) from highly complex sample matrix.

Acknowledgements: T. C. F. Ribas and R. B. R. Mesquita thank to Fundação para a Ciência e a Tecnologia (FCT, Portugal) and POCH of Fundo Social Europeu (FSE) for the grants SFRH/BD/91820/2012 and SFRH/BPD/112032/2015, respectively. This work was also supported by National Funds from FCT - Fundação para a Ciência e a Tecnologia through the project PTDC/AAG-MAA/5887/2014; scientific collaboration from FCT project UID/Multi/50016/2013 is also acknowledged.

[1] Fontàs C., Tayeb R., Dhahbi M., Gaudichet E., Thominet F., Roy P., Steenkeste K., Fontaine-Aupart M., Tingry S. *J. Membr. Sci.* **2007**, *290*, 62-72.

[2] Nghiem L. D., Mornane P., Potter I. D., Perera J. M., Cattrall R. W., Kolev S. D. *J. Membr. Sci.* **2006**, *281*, 7-41.

TÍTULO

Book of Abstracts Analítica 2018 - 9th Meeting
of Division of Analytical Chemistry

EDITORIAL BOARD

Marcela A. Segundo
Eduarda M. P. Silva
Luísa Barreiros
Lúcia Saraiva

TYPESET AND GRAPHIC DESIGN

Joana Macedo, FFUP

ALL RIGHTS RESERVED

This publication may not be reproduced or
transmitted in whole or in part by any process,
electronic, mechanical, photocopying, recording
or otherwise, without the prior permission of the
editorial board.

EDITION

Sociedade Portuguesa de Química
Av. da República, 45 - 3^o Esq
1050-187 Lisboa - Portugal
ISBN: 978-989-8124-21-0

DATE

March 2018